

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>A61K 47/44, 31/445</b>		<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/32234</b>
			<b>(43) International Publication Date:</b> 8 June 2000 (08.06.00)
<b>(21) International Application Number:</b> PCT/EP99/09351 <b>(22) International Filing Date:</b> 1 December 1999 (01.12.99)  <b>(30) Priority Data:</b> 9826656.2      3 December 1998 (03.12.98)      GB  <b>(71) Applicant (for all designated States except AT US):</b> NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).  <b>(71) Applicant (for AT only):</b> NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT MBH . [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> KRIWET, Katrin [DE/DE]; Allmendweg 9, D-79639 Grenzach-Wyhlen (DE). LEDERGERBER, Dorothea [DE/DE]; Brunnenweg 11a, D-79539 Lörrach (DE). RIEDL, Jutta [DE/DE]; Rebgasse 17, D-79639 Grenzach (DE).  <b>(74) Agent:</b> BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Department, CH-4002 Basel (CH).			<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> TOPICAL COMPOSITIONS COMPRISING ASCOMYCINS			
<b>(57) Abstract</b>  The present invention relates to a composition for topical administration comprising an ascomycin and a carrier vehicle comprising means to retain water in the outer skin layer and means to hinder water evaporating from the skin.			
<b>BEST AVAILABLE COPY</b>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## TOPICAL COMPOSITIONS COMPRISING ASCOMYCINS

This invention relates to topical compositions containing ascomycins for treatment of skin disorders, e.g. subacute and chronic inflammatory and hyperproliferative skin diseases, e.g. atopic dermatitis, vitiligo, psoriasis, lichenified skin diseases, e.g. lichen planus, a lichenified  
5 form of atopic dermatitis.

Ascomycins have a variety of useful pharmacological actions, e.g. immunosuppression, and which may be administered topically. However, inter alia because of their physicochemical properties, e.g. high molecular weight and lipophilicity the ascomycins have posed problems  
10 for topical administration.

Skin disorders also present difficulties in treatment, particularly lichenified skin diseases, e.g. psoriasis, where the skin is hyperproliferated and the skin barrier function and skin lipid composition may have changed. Topical compositions for use in lichenified skin diseases, e.g. psoriasis, containing an ascomycin present particular difficulties.  
15

After exhaustive testing it has now been surprisingly found that the compositions of the present invention serve to enhance penetration of active agent through human skin, e.g. for the treatment of lichenified skin diseases, e.g. psoriasis. These compositions show other particularly interesting properties, e.g. they are easily applied to large areas of the skin and  
20 are stable.

In one aspect this invention provides a composition for topical administration of an ascomycin which composition comprises a carrier vehicle comprising

- (i) means to retain water in the outer skin layer, and
- 25 (ii) means to hinder water evaporating from the skin.

The ascomycin is hereafter referred to as active agent. Under "ascomycin" is to be understood ascomycin itself or a derivative, antagonist, agonist or analogue thereof, e.g. a compound of the FK 506 class.

30 FK506 is a known macrolide antibiotic that is produced by Streptomyces tsukubaensis No 9993. It is also a potent immunosuppressant. The structure of FK506 is given in the

appendix to the Merck Index, 11th Edition as item A5. Methods of preparing FK506 are described in EP 184162.

Under "compound of the FK 506 class" is to be understood FK 506 itself or a derivative,  
5 antagonist, agonist or analogue thereof, which retain the basic structure and modulate at least one of the biological properties (for example immunological properties) of FK506. A large number of compounds of the FK 506 class are known. These compounds are described in for example EP 184162, EP 315978, EP 323042, EP 423714, EP 427680, EP 465426, EP 474126, WO 91/13889, WO 91/19495, EP 484936, EP 532088, EP 532089,  
10 EP 569337, EP 626385, WO 93/5059 and the like.

It is also known (for example from EP 315978 and EP 474126) that ascomycin derivatives such as macrolactam compounds of the FK506 class are particularly useful in the topical treatment of inflammatory and hyperproliferative skin diseases and of cutaneous  
15 manifestations of immunologically-mediated illnesses.

Thus examples of ascomycin derivatives suitable for use in the present invention include FK506; 33-epi-chloro-33-desoxy-ascomycin as disclosed in Example 66a in EP 427 680 (hereafter referred to as Compound A);

20

{[1E-(1R,3R,4R)]1R,4S,5R,6S,9R,10E,13S,15S,16R,17S,19S,20S}-9-ethyl-  
6,16,20-trihydroxy-4-[2-(4-hydroxy-3-methoxy-cyclohexyl)-1-methylvinyl]-  
15,17-dimethoxy-5,11,13,19-tetramethyl-3-oxa-22-aza-tricyclo[18.6.1.0(1,22)]

heptacos-10-ene-2,8,21,27-tetraone as disclosed in Examples 6d and 71 in EP 569 337

25 (hereafter referred to as Compound B); and

{1R,5Z,9S,12S-[1E-(1R,3R,4R)],13R,14S,17R,18E,21S,23S,24R,25S,27R}  
17-ethyl-1,14-dihydroxy-12-[2-(4-hydroxy-3-methoxy-cyclohexyl)-1-methylvinyl]  
-23,25-dimethoxy-13,19,21,27-tetramethyl-11,28-dioxa-4-aza-tricyclo[22.3.1.0(4,9)]

30 octacos-5,18-diene-2,3,10,16-tetraone, also known as 5,6-dehydro-ascomycin as disclosed in Example 8 in EP 626 385 (hereafter referred to as Compound C);

Imidazolylmethoxyascomycin, as disclosed in Example 1 and as compound of formula I in WO 97/08182, contents of which are incorporated herein by reference (hereafter referred to as Compound D);

- 5 32-O-(1-hydroxyethylindol-5-yl)ascomycin, also known as Indolyl-ASC or L-732 531 as disclosed in Transplantation 65 (1998) 10-18, 18-26, on page 11, Figure 1 (hereafter referred to as Compound E); and

- 10 (32-deoxy-32-epi-N1-tetrazolyl)ascomycin, also known as ABT-281 as disclosed in J. Inv. Derm. 112 (May 1999), 729-738, on page 730, Figure 1 (hereafter referred to as Compound F).

FK 506, Compounds A, B, C, D, E, and F are preferred ascomycins, particularly preferred are Compounds A, B, and C, especially Compound A.

15

The active agent is e.g. present in the compositions of this invention in an amount of from 0.05 to 3 % by weight, e.g. from 0.1 to 2 % by weight, e.g. from 0.4 to 1 % by weight based on the total weight of the composition.

- 20 The active agent may be dissolved, e.g. partially dissolved in the vehicle. In a further aspect the active agent may be in suspension, e.g. partially in suspension in the vehicle. Preferably the active agent is partially dissolved in the vehicle.

- 25 Preferably the active agent may be used in a micronized form. The suspension may contain particles of ascomycin of from 5, e.g. from 10, to about 90, preferably to about 25 microns in diameter. The particles of the ascomycin may be produced in conventional manner, e.g. by grinding or milling.

If desired further active agents may be present.

- 30 The carrier vehicle comprises means to retain water in the outer skin layer, e.g. moisturizers.

Under "means to retain water in the outer skin layer" is to be understood, e.g. a pharmaceutically acceptable moisturizer, capable of e.g. penetrating and residing in the

outer skin layer, e.g. the stratum comeum, and e.g. absorbing, holding and retaining moisture to increase the moisture content of the skin.

Means to retain water in the outer skin layer, e.g. moisturizers, e.g. as described in

5 Dermatika, Eds. R. Nieder, J. Ziegenmayer, Wissenschaftliche Verlagsgesellschaft Stuttgart 1992, 271-272, may be selected from a group comprising

- i) a urea, e.g. urea and its derivatives, e.g. monoacetyl urea, 1-dodecyl urea, 1,3-didodecyl urea, 1,3 diphenyl urea or cyclic urea derivatives, e.g. 1-methyl-4-imidazolin-2-one-3-methylenedecanoate. Urea may be commercially available from  
10 e.g. Merck, Germany;
- ii) an inorganic salt, e.g. sodium chloride, e.g. as known and commercially available from e.g. Merck, Germany; and
- iii) a carboxylic acid, e.g. a mono carboxylic acid or a cyclic carboxylic acid, salts and  
15 derivatives thereof. Particularly preferred are e.g. lactic acid; glycolic acid; lactic acid sodium and/or ammonium salt, e.g. sodium lactate, e.g. as known and commercially available from e.g. Merck, Germany; glycolic acid sodium and/or ammonium salt; lactamide; lactamidopropyl-triammonium chloride; sodium cocoyl lactylate; 2-pyrrolidone-5- carboxylate; 2-pyrrolidone-5- carboxylate sodium and/or calcium salt,  
20 e.g. sodium 2-pyrrolidone-5-carboxylate, e.g. as known and commercially available under the name Sodium PCA from A+E Connock, UK; 2-pyrrolidone-5- carboxylate derivatives of amino acids, e.g. lysin or arginin; or acyl esters, having e.g. a chain length of C<sub>1</sub>-C<sub>30</sub>, e.g. C<sub>3</sub>-C<sub>18</sub>, branched or unbranched, e.g. 2-pyrrolidone-5- carboxylic laurate (Fiedler, H.P. "Lexikon der Hilfsstoffe für Pharmazie, Kosmetik und angrenzende Gebiete", Editio Cantor Verlag Aulendorf, Aulendorf, 4th revised and  
25 expanded edition (1996), 1, p. 720; 2, p. 1013-1017; 2, p. 1303)

Mixtures thereof may also be used.

Preferably the carrier vehicle comprises a urea, e.g. urea itself or derivatives thereof, e.g. monoacetyl urea, 1-dodecyl urea, 1,3- didodecyl urea, 1,3 diphenyl urea or cyclic urea  
30 derivatives, e.g. 1-methyl-4-imidazolin-2-one-3-methylenedecanoate.

Particularly preferred is urea itself.

The means to retain water in the outer skin layer, e.g. urea, may be present in amount of from 0.1 to about 20 %, e.g. from 1 to about 15 %, preferably about 5 % by weight based

on the total weight of the composition. The means to retain water in the outer skin layer may be suspended or dispersed in the vehicle. They may be used in a micronized or non micronized form. Particularly preferred is the micronized form. The suspension or dispersion may contain particles of, e.g., urea of from 5, e.g. from 10, to about 90, preferably to about 25 microns in diameter. The non-micronized particles may have a size of equal to or less than 500 microns. The particles of the urea may be produced in conventional manner, e.g. by grinding or milling.

Preferably the ascomycin and the means to retain water in the outer skin layer are present in a weight ratio of 0.05 to 3 : 0.1 to 20, more preferably in a weight ratio of 0.1 to 2 : 5 to 15, even more preferably in a weight ratio of 0.4 to 1 : about 5.

The carrier vehicle further comprises means to hinder water evaporating from the skin, e.g. hydrocarbons. Hydrocarbons may be selected from a group comprising

- i) petrolatum, e.g. white petrolatum, e.g. as known and commercially available from e.g. Mineral Chemie AG, Germany;
- ii) liquid paraffin, e.g. as known and commercially available from e.g. Mobil BP Oiltech, Switzerland;
- iii) solid paraffin; or microcrystalline wax, e.g. as known and commercially available under the trade name Esma® M from Schlüter, Germany; and
- iv) a reaction product of a paraffin and a polyethylene, e.g. a polyethylene having a molecular weight of from 10000 to about 400000 Daltons, e.g. 21000 Daltons, e.g. as known under the name Hydrophobes Basisgel DAC and commercially available under the trade name Plastibase®, from e.g. Hansen & Rosenthal, Germany (Fiedler, H.P., loc. cit., 2, p. 1198).

Mixtures thereof may also be used.

Hydrocarbons may be present in amount of from 70 to about 95 %, preferably of from 75 to about 90 %, more preferably about 85 % by weight based on the total weight of the composition.

The amount and the type of hydrocarbons in the composition may depend on the desired viscosity of the composition as is conventional.

Preferably the ascomycin and the hydrocarbon are present in a weight ratio of 0.05 to 3 : 70 to 95, more preferably in a weight ratio of 0.1 to 2 : 75 to 90, even more preferably in a weight ratio of 0.4 to 1 : about 85.

5 In another aspect the present invention provides a composition as defined above which composition comprises a carrier vehicle comprising

- (i) a urea, an inorganic salt, or a carboxylic acid, and
- (ii) a hydrocarbon.

10 Under "a carboxylic acid" is to be understood a mono carboxylic acid or a cyclic carboxylic acid, salts and derivatives thereof, e.g. as defined above. Under "a urea" is to be understood urea it self or a derivative thereof, e.g. as defined above.

In another aspect the present invention provides a composition as defined above which

15 composition comprises a carrier vehicle further comprising

- (iii) liquid means, e.g. lipophilic solvents and/or polar solvents, to solubilize ascomycin.

The lipophilic solvents may be selected from a group comprising

- 20 i) liquid waxes, e.g. natural-, synthetic-, semisynthetic- or emulsifying- waxes. Preferably isopropyl myristate, e.g. as known and commercially available from Henkel, Germany; oleyl erucate, e.g. as known and commercially available under the trade name Cetiol® J600 from e.g. Henkel, Germany; diisopropyl adipate, e.g. as known and commercially available under the trade name Isopat® 1794 from e.g. Dargoco, Germany; and/or oleyl oleate, e.g. as known and commercially available under the trade name Cetiol®  
25 from e.g. Henkel, Germany, may be used;
- ii) liquid fatty alcohols, saturated and/or unsaturated, branched and/or unbranched, having e.g. a C<sub>8</sub> to C<sub>24</sub> chain. Preferably oleyl alcohol, e.g. as known and commercially available under the trade name HD Eutanol® from e.g. Henkel, Germany, may be used;
- 30 iii) fatty acids, saturated and/or unsaturated, branched and/or unbranched, having e.g. a C<sub>8</sub> to C<sub>24</sub> chain, e.g. oleic acid and/or lauric acid; and
- iv) fatty oils, comprising e.g. mono-, di- and tri- glycerides, having e.g. C<sub>8</sub> to C<sub>24</sub> fatty acids, e.g. a medium chain fatty acid triglyceride, e.g. Miglyol®812. Miglyol®812 is a



fractionated coconut oil comprising caprylic-capric acid triglycerides and having a molecular weight of about 520 daltons. Fatty acid composition = C<sub>6</sub> max. about 3%, C<sub>8</sub> about 50 to 65%, C<sub>10</sub> about 30 to 45%, C<sub>12</sub> max 5%; acid value about 0.1; saponification value about 330 to 345; iodine value max 1. Miglyol® 812 is commercially available from e.g. Hüls Chemie AG, Germany.

The polar solvents may be selected from a group comprising

- i) glycols, e.g. glycerol, propylene glycol, butylene glycol, hexylene glycol. Propylene glycol may be commercially available from e.g. Dow Chemical;
- ii) alcohols, having e.g. a C<sub>1</sub> to C<sub>7</sub> chain, branched and/or unbranched, e.g. isopropanol;
- 10 iii) dimethyl isosorbide, e.g. as known and commercially available under the trade name Arlasolve® DMI from ICI, Germany; and
- iv) propylene carbonate.

The liquid means to solubilize the ascomycin may consist of one component or a mixture of components. Preferably the liquid means may be isopropyl myristate. The liquid means may be present in amount of from 1 to 20 %, preferably from 2 to 15 %, more preferably about 5 % by weight based on the total weight of the composition.

Preferably the composition is in the form of an ointment, containing no added water, e.g. a water content of less than 5 or 2%.

20

The liquid means may serve to dissolve partially the active agent. Typically 1 to 5 % of the active agent is dissolved. Preferably a saturated solution of the active agent in the composition is obtained.

25 Preferably the ascomycin and the liquid means are present in a weight ratio of 0.05 to 3 : 1 to 15, more preferably in a weight ratio of 0.1 to 2 : 2 to 10, even more preferably in a weight ratio of 0.4 to 1 : about 5.

30 Preferably the ascomycin, the urea, the hydrocarbon and the liquid means, when present, are present in a weight ratio of 0.05 to 3 : 0.1 to 20 : 70 to 95 : 1 to 15, more preferably in a weight ratio of 0.1 to 2 : 5 to 15 : 75 to 90 : 2 to 10, even more preferably in a weight ratio of 0.4 to 1 : about 5 : about 85 : about 5.

The components of the carrier vehicle may be described in Fiedler, H.P., loc. cit., the contents of which are hereby incorporated by reference.

5 The compositions of this invention may be water-free or substantially water-free. The compositions may however comprise water, e.g. in an amount of from 0 to about 10 % by weight based on the total weight of the composition, e.g. from 0.5 to 5 %, e.g. from 1 to 3 %. Preferably the compositions of this invention may be water-free.

The compositions of the invention are preferably in the form of an ointment.

10 If desired, stabiliser agents to hinder degradation of urea may be included, e.g. allantoin, acetyl glyceride, propionic acid ester, taurin, collagen, collagen hydrolysate, amino acid salts, monoalkylphosphate diethanolamine, triacetin, lactic acid, polysaccharides, chelating agents, e.g. citric acid or EDTA, e.g. as described in Fiedler, H.P. (loc. cit., 1, page 737).

15 Further components, e.g. preserving agents, e.g. microorganism growth inhibitors, and antioxidants, such as benzyl alcohol, butyl-hydroxytoluene, ascorbyl palmitate, sodium pyrosulfite, butyl hydroxy anisole, propyl p-hydroxybenzoate, methyl p-hydroxybenzoate, sorbic acid, chlorcresol and tocopherol may be included as appropriate. Preserving agents  
20 and antioxidants are preferably present in an amount of about 0.01 to about 2.5 % by weight based on the total weight of the composition.

If desired, pH modifying agents may be included to bring the pH of the composition to between 4 and 6 or by adding a pharmaceutically acceptable buffer system. A pH of  
25 between 4 and 6 is desirable to avoid skin irritation.

If desired, the compositions of the invention may further comprise thickeners, e.g. to stabilize the compositions, e.g.

- 30 i) solid alcohols, having e.g. a C<sub>12</sub> to C<sub>24</sub> chain, e.g. cetyl alcohol and/or stearyl alcohol. Cetyl alcohol and stearyl alcohol may be commercially available e.g. under the trade names Lorol® C16 and Lorol® C18, respectively, from Henkel, Germany;
- ii) solid acids, having e.g. a C<sub>12</sub> to C<sub>24</sub> chain, e.g. stearic acid and its salts, e.g. aluminium- or magnesium stearate;

- iii) esters, e.g. solid esters, of glycerol, e.g. mono-, di-, or tri- esters, e.g. glycerol monostearate and/or hydrogenated castor oil. Glycerol monostearate may be commercially available under the trade name Atmul® 84K from ICI, Germany;
- iv) esters, e.g. solid esters, of propylene glycol, e.g. mono- or di- esters, e.g. propylene glycol monooleate;
- 5 v) inorganic thickening agents, e.g. magnesium sulfate, bentonite or silicates including hydrophilic silicon dioxide products, e.g. alkylated, for example methylated, silica gels, in particular colloidal silicon dioxide products as known and commercially available under the trade name Aerosil, e.g. Arosil® 200, Aerosil® R812 or Aerosil® R 972, e.g.
- 10 from Degussa, Germany (Handbook of Pharmaceutical Excipients, 2nd Edition, Editors A. Wade and P. J. Weller (1994), Joint publication of American Pharmaceutical Association, Washington, USA and The Pharmaceutical Press, London, England, page 424-427);
- vi) solid waxes, e.g. bees wax or carnauba wax; and
- 15 vii) esterified compounds of fatty acid and fatty alcohols. They may include esterified compounds of fatty acid having e.g. a C<sub>12</sub> to C<sub>24</sub> chain, saturated or unsaturated, and primary alcohol having e.g. a C<sub>12</sub> to C<sub>24</sub> chain, e.g. cetyl palmitate.

Thickening agents are preferably present in an amount of from about 1 % to about 30 %, e.g. from about 2 % to about 10 %, by weight based on the total weight of the composition.

20

The compositions of the present invention may further comprise emulsifiers, e.g.

- i) Polyoxyethylene-sorbitan-fatty acid esters, for example mono- and tri-lauryl, palmityl, stearyl and oleyl esters of the type known and commercially available under the trade name Tween® (Fiedler, loc.cit. p.1615 ff), including the products Tween®
- 25 20 [polyoxyethylene(20)sorbitanmonolaurate],  
21 [polyoxyethylene(4)sorbitanmonolaurate],  
40 [polyoxyethylene(20)sorbitanmonopalmitate],  
60 [polyoxyethylene(20)sorbitanmonostearate],  
65 [polyoxyethylene(20)sorbitantristearate],  
80 [polyoxyethylene(20)sorbitanmonooleate],  
30 81 [polyoxyethylene(5)sorbitanmonooleate],

85 [polyoxyethylene(20)sorbitantrioleate].

Especially preferred products of this class are Tween® 60 and Tween® 65.

- 5      ii) Sorbitan fatty acid esters, e.g. sorbitan mono C<sub>12-18</sub> fatty acid esters, or sorbitan tri C<sub>12-18</sub> fatty acid esters as known and commercially available under the trade mark Span® or Arlacel®. Particularly preferred are the products Arlacel® 83 (Sorbitan sesquioleate) available from ICI, Germany, or Span® 60 (Sorbitan monostearate) (Fiedler, loc. cit., 2, p. 1430; Handbook of Pharmaceutical Excipients, loc. cit., page 473).
- 10      iii) Polyoxyethylene alkyl ethers, e.g. polyoxyethylene glycol ethers of C<sub>12</sub> to C<sub>18</sub> alcohols, e.g. Polyoxyl 2-, 10- or 20-cetyl ether or Polyoxyl 4- or 23-lauryl ether, or polyoxyl 2-, 10- or 20-oleyl ether, or Polyoxyl 2-, 10-, 20- or 100-stearyl ether, as known and commercially available under the trade name Brij® from e.g. ICI, Germany. An especially preferred product of this class is e.g. Brij® 30 (Polyoxyl 4 lauryl ether) or
- 15      Brij® 72 (Polyoxyl 2 stearyl ether) (Fiedler, loc. cit., 1, pp. 259; Handbook of Pharmaceutical Excipients, loc. cit., page 367).
- 20      iv) Polyoxyethylene fatty acid esters, for example polyoxyethylene stearic acid esters of the type known and commercially available under the trade name Myrj® (Fiedler, loc. cit., 2, p. 1042; Handbook of Pharmaceutical Excipients, loc. cit., page 379). An especially preferred product of this class is Myrj® 52 (Polyoxyethylene 40 stearate) having a D<sup>25</sup> of about 1.1., a melting point of about 40 to 44°C, an HLB value of about 16.9., an acid value of about 0 to 1 and a saponification no. of about 25 to 35.
- 25      v) Sucrose esters, e.g. sucrose fatty acid esters. The fatty acid moiety may comprise saturated or unsaturated fatty acids or mixtures thereof. Particularly suitable are C<sub>6-18</sub> fatty acid saccharide mono- or diesters, in particular water soluble C<sub>6-18</sub> fatty acid saccharide mono- or diesters. Especially suitable are caproic (C<sub>6</sub>), caprylic (C<sub>8</sub>), capric (C<sub>10</sub>), lauric (C<sub>12</sub>), myristic (C<sub>14</sub>), palmitic (C<sub>16</sub>), oleic (C<sub>18</sub>), ricinoleic (C<sub>18</sub>) and 12-
- 30      hydroxystearic (C<sub>18</sub>) acid saccharide mono- or diesters, e.g. sucrose distearate, e.g. as known and commercially available under the trade name Sucro Ester® 7 from Gattefossé, France.

vi) Silicone emulsifiers, e.g. laurylmethicone copolyol, e.g. as known and commercially available under the trade name Emulsifier® 10 from Dow Corning or a mixture of cetyldimethicone copolyol, polyglyceryl-4-isostearate and hexyl laurate, e.g. as known and commercially available under the trade name Abil® WE-09 from Goldschmidt.

5

vi) Phospholipids, in particular lecithins (Fiedler, H. P., "Lexikon der Hilfsstoffe für Pharmazie, Kosmetik und angrenzende Gebiete", Editio Cantor Verlag Aulendorf, Aulendorf, 4th revised and expanded edition (1996), vol 2, p. 910, 1184). Lecithins suitable for use in the compositions of the invention include egg lecithins or soybean lecithins, in particular soybean lecithins, e.g. as known and commercially available under the trade name Phospholipon® 80 from Rhone Poulenc Rorer. Phospholipon® 80 is a phospholipid fraction with about 76 % phosphatidylcholine, about 8 % phosphatidic acid, about 4 % phosphatidyl ethanolamine, and about 9 % other lipids (manufacturer information).

10

15

vii) Lanolin, e.g. anhydrous lanolin (Fiedler, H.P., loc.cit., 2, p. 896).

It is to be appreciated that emulsifiers may be complex mixtures containing side products or unreacted starting products involved in the preparation thereof, e.g. emulsifiers made by polyoxyethylation may contain another side product, e.g. polyethylene glycol.

20

Compositions additionally comprising emulsifiers may be particularly suitable if it is desirable to easily wash them off the skin.

25 The compositions of the invention may further include, e.g. perfumes and/or coloring agents, as appropriate.

The compositions according to the invention are useful in the treatment of subacute and chronic inflammatory and hyperproliferative skin diseases and of cutaneous manifestations of immunologically-mediated diseases. Examples of such diseases are psoriasis, atopic dermatitis, contact dermatitis and further eczematous dermatitises, seborrheic dermatitis, Lichen planus, a lichenified form of atopic dermatitis, vitiligo, Pemphigus, bullous

30

Pemphigoid, Epidermolysis bullosa, urticaria, angioedemas, vasculitides, erythemas, cutaneous eosinophilias, Lupus erythematosus and Alopecia areata.

5 In another aspect the present invention provides a composition as defined above for use in the treatment of inflammatory and hyperproliferative skin diseases and of cutaneous manifestations of immunologically-mediated diseases.

10 In another aspect the present invention provides a method for treating inflammatory and hyperproliferative skin diseases and of cutaneous manifestations of immunologically-mediated diseases comprising administering a composition as defined above to the skin of a patient in need thereof.

15 In another aspect the present invention provides the use of a composition as defined above in the preparation of a medicament for administering to the skin of a patient in need thereof.

20 In yet another aspect the present invention provides the use of a composition as defined above in the preparation of a medicament for the treatment of inflammatory and hyperproliferative skin diseases and of cutaneous manifestations of immunologically-mediated diseases.

25 In yet another aspect the present invention provides the use of a carrier vehicle as defined above to enhance penetration of an ascomycin through human skin.

30 The carrier vehicle may be in the form of an ointment.

The compositions of the invention may be prepared in a conventional manner by working up the components into a pharmaceutical composition.

For example, the composition of the invention may be obtained by suspending the ascomycin and the urea in a mixture of liquid hydrocarbons and the lipophilic or polar solvent. Solid hydrocarbons may be mixed into the suspension in conventional manner. Alternatively, the composition of the invention may be obtained by suspending the ascomycin and the urea in a mixture of liquid hydrocarbons, solid hydrocarbons and the solvent as conventional. Other, e.g. conventional, excipients may be added at the appropriate time.

Claims

1. A composition for topical administration of an ascomycin which composition comprises a carrier vehicle comprising
  - 5 (i) means to retain water in the outer skin layer, and
  - (ii) means to hinder water evaporating from the skin.
2. A composition as claimed in claim 1 wherein the means to retain water in the outer skin layer comprises a urea, an inorganic salt, or a carboxylic acid.
- 10 3. A composition as claimed in claim 2 wherein the means to retain water in the outer skin layer is a urea.
4. A composition as claimed in any one of claims 1 to 3 wherein the means to hinder water evaporating from the skin is a hydrocarbon.
- 15 5. A composition as claimed in claim 4 wherein the hydrocarbon comprises petrolatum, liquid paraffin, microcrystalline wax, solid paraffin, or a reaction product of paraffin and polyethylene.
- 20 6. A composition as claimed in any preceding claim wherein the carrier vehicle further comprises
  - (iii) liquid means to solubilize ascomycin.
- 25 7. A composition as claimed in claim 6 wherein the liquid means comprises a wax, a fatty alcohol, a fatty acid, or a fatty oil.
8. A composition as claimed in claim 7 wherein the liquid means is isopropyl myristate.
- 30 9. A composition as claimed in any preceding claim wherein the ascomycin is present in an amount of 0.1 to 2.0 % by weight of the composition.

# INTERNATIONAL SEARCH REPORT

information on patent family members

Info

PCT

Patent document cited in search report		Publication date	Patent family member(s)	
EP 0423714	A	24-04-1991	AT	107499 T
			CA	2027608 A
			DE	69010139 D
			DE	69010139 T
			DK	423714 T
			JP	2925285 B
			JP	3204807 A
			KR	159766 B
			US	5215995 A
EP 0474126	A	11-03-1992	AT	150304 T
			AU	656145 B
			AU	8351591 A
			CA	2050623 A
			CN	1059468 A
			DE	69125230 D
			DE	69125230 T
			DK	474126 T
			ES	2099112 T
			GR	3022883 T
			HK	1000006 A
			HU	59002 A
			JP	2526752 B
			JP	5017481 A
			PT	98862 A
			SG	46547 A
			RU	2079303 C
			US	5385907 A
			ZA	9106983 A
WO 9613249	A	09-05-1996	AU	714254 B
			AU	3845195 A
			AU	5833699 A
			BR	9509530 A
			CA	2200966 A
			CZ	9701232 A
			DE	19581804 T
			EP	0786986 A
			FI	971018 A
			GB	2308546 A
			HU	77140 A
			JP	10508588 T
			NO	971951 A
			NZ	295170 A
			PL	319599 A
			SK	52097 A
			GB	2327610 A

BEST AVAILABLE COPY



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**